arXiv:0905.2041v3 [astro-ph.CO] 15 Jun 2010

Scalar field fluctuations in Schwarzschild-de Sitter space-time

Hing-Tong Cho[1], Kin-Wang Ng[2], and I-Chin Wang[3]

[1] *Department of Physics, Tamkang University, Tamsui, Taiwan 251, R.O.C.*
[2] *Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.*
[3] *Department of Physics, National Taiwan Normal University, Taipei, Taiwan 116, R.O.C.*

(Dated: June 16, 2010)

Abstract

We calculate quantum fluctuations of a free scalar field in the Schwarzschild-de Sitter space-time, adopting the planar metric that is pertinent to the presence of a black hole in an inflationary universe. In a perturbation approach, doing expansion in powers of a small black hole mass, we obtain time evolution of the quantum fluctuations and then derive the scalar power spectrum.

PACS numbers: 04.62.+v, 04.70.Bw, 98.80.Cq

I. INTRODUCTION

The cosmic microwave background that we observe today is almost homogenous and isotropic. The background temperature in our sky is about 2.7K with a tiny fluctuation at a level of about 10^{-5}K. This is consistent with measurements of the matter content of the Universe, altogether prevailing a spatially flat universe. Inflation scenario can explain the homogeneity, the isotropy, and the flatness of the present Universe. Moreover, quantum fluctuations of the inflaton field during inflation can give rise to primordial density fluctuations with a nearly scale-invariant power spectrum which is consistent with the recent WMAP data on cosmic microwave background anisotropies [1]. Therefore, a general assumption usually made in most cosmological models is that the background metric is homogenous and isotropic. For example, a de Sitter metric is used in the inflationary era and a flat Friedmann-Robertson-Walker metric is used in the subsequent hot big bang. An interesting notion is the recent discovery of a dominant component in the matter content, dubbed the dark energy, which exerts a negative pressure to drive an accelerated expansion of the Universe. If dark energy is a form of vacuum energy, our Universe will coast to the de Sitter space-time or the inflating phase again in the future.

Apparently our present Universe is not so homogeneous and isotropic because we observe local non-linear structures such as stars, galaxies, clusters of galaxies, and very massive black holes. An appropriate space-time, for example, for a massive black hole sitting in the accelerating Universe, would be described by the Schwarzschild metric in the vicinity of the black hole and by the de Sitter metric at places far from the black hole. Presently, these local structures are decoupled from the Hubble flow, so it suffices to use the Friedmann-Robertson-Walker metric to study the large scale structures of the Universe. However, in the early Universe the gravitational effect of a black hole to the background metric may be important and should be addressed. For example, the existence of a black hole or a distribution of black holes at the onset of inflation can invalidate the use of a homogenous and isotropic background metric for the calculation of de Sitter quantum fluctuations. This also applies to the situation when we are very near to one of these black holes that still exists today or not.

The cosmic no hair conjecture infers that the inflationary universe approaches asymptotically the de Sitter space-time till the end of inflation [2]. Nevertheless, the effects of matter and space-time inhomogeneities to inflation should be considered. Several authors have studied the onset of inflation under inhomogeneous initial conditions to determine whether large inhomogeneity during the very early Universe can prevent the Universe from entering an inflationary era [3]. It was found that in some cases a large initial inhomogeneity may suppress the onset of inflation [4]. If the inflaton field is sufficiently inhomogeneous, the wormhole can form before the inhomogeneity is damped by the exponential expansion [5]. In the case of inhomogeneities in a dust era before inflation, some inhomogeneities can collapse into a black-hole space-time [6]. Furthermore, for the inhomogeneities of the space-time itself, energies in the form of gravitational waves can also form a black-hole space-time [7]. As a consequence, at the onset of inflation, the distortion of the metric by these inhomogeneities should be taken into account.

With these considerations in mind, in this work we will investigate the quantum fluctuations of a free massless scalar field in the Schwarzschild-de Sitter (SdS) space-time. In the static coordinate system, the line element of the SdS space-time is given by

$$ds^2 = -\left(1 - \frac{2GM}{r} - H^2r^2\right)dt^2 + \left(1 - \frac{2GM}{r} - H^2r^2\right)^{-1} dr^2 + r^2d\Omega^2, \tag{1}$$

where $G = M_{\rm Pl}^2$, M is the mass of the black hole, and H is the Hubble parameter for inflation. Here we use the convention with $c = \hbar = 1$. As is well-known, the SdS metric has a black hole horizon and a cosmological horizon. In the static coordinates (1) an observer can only receive a signal inside or just right on the cosmological horizon. This static metric is insufficient for our purpose because in the cosmological setting we aim at studying the temporal evolution of a Fourier mode of the scalar quantum fluctuations that crosses the cosmological horizon during inflation. Therefore, we will instead use the planar coordinates for the SdS metric [8], which is given by

$$ds^2 = -f(r,\tau)d\tau^2 + h(r,\tau)(dr^2 + r^2d\Omega^2), \tag{2}$$

where $d\tau = a^{-1}(\tau)dt$ is the conformal time and $a(t) = e^{Ht}$. In Eq. (2), the f and h functions are given by

$$f(r,\tau) = a^2(\tau)\left[1 - \frac{GM}{2a(\tau)r}\right]^2\left[1 + \frac{GM}{2a(\tau)r}\right]^{-2}, \quad h(r,\tau) = a^2(\tau)\left[1 + \frac{GM}{2a(\tau)r}\right]^4, \tag{3}$$

with the cosmic scale factor $a(\tau) = -1/(H\tau)$. Note that at large times the planar coordinates behave like a de Sitter expansion. Also, we can cast it to a co-moving frame that is conformal to the Minkowskian space-time with a black hole in it.

It is well-known that a black hole evaporates into Hawking radiation which leads to a mass loss of the black hole and to its eventual disappearance [9]. For a Schwarzschild black hole with mass M, the evaporation time is given by

$$t_{\rm ev} = 5120\pi G^2M^3. \tag{4}$$

Therefore, our present consideration requires the condition that the evaporation time scale of the black hole is longer than the time scale of inflation, i.e., $Ht_{\rm ev} > 1$. This gives the lower bound on the mass of the black hole for a given inflation scale:

$$\frac{M}{M_{\rm Pl}} > 3.96 \times 10^{-2}\left(\frac{M_{\rm Pl}}{H}\right)^{\frac{1}{3}}. \tag{5}$$

In the next section, we will review the scalar quantum fluctuations in the de Sitter metric. Then, we will introduce a perturbation method to expand the planar metric in powers of the black hole mass M to find an approximate solution of the scalar equation. Section III contains the numerical results of the first-order solutions. Section IV is our conclusion.

II. PERTURBATION APPROACH TO THE SCHWARZSCHILD-DE SITTER SPACE-TIME

A. Classical solution

Consider a massless scalar field ϕ which satisfies the Klein-Gordon equation in the SdS space-time,

$$\partial_\mu\left(\sqrt{-g}g^{\mu\nu}\partial_\nu\phi\right) = 0. \tag{6}$$

Since the space is spherically symmetric, one can expand ϕ as

$$\phi(x) = \sum_{lm} \varphi_l(r,\tau) Y_{lm}(\theta,\phi). \tag{7}$$

If one further writes $\varphi_l(r,\tau)$ in a spectral form,

$$\varphi_l(r,\tau) = \int_0^\infty dk\, k^2 j_l(kr) \varphi_{kl}(\tau), \tag{8}$$

then we will have

$$\phi(x) = \int_0^\infty dk \sum_{lm} \varphi_{klm}(x), \quad \varphi_{klm}(x) = k^2 j_l(kr)\varphi_{kl}(\tau) Y_{lm}(\theta,\phi). \tag{9}$$

We define the spectral function of the fluctuations of the field $\phi(x)$ as

$$S_{kl}(r,\tau) \equiv (2l+1) j_l^2(kr) P_{kl}(\tau), \quad P_{kl}(\tau) = \frac{k^5}{4\pi}|\varphi_{kl}(\tau)|^2. \tag{10}$$

Our next task is to calculate the function $\varphi_{kl}(\tau)$ in Eq. (10) in the SdS planar metric (2). The Klein-Gordon equation (6) beomes

$$-\frac{1}{\sqrt{fh}}\partial_\tau\left(\sqrt{\frac{h^3}{f}}\partial_\tau\varphi_l\right) + \frac{1}{r^2\sqrt{fh}}\partial_r\left(r^2\sqrt{fh}\partial_r\varphi_l\right) - \frac{l(l+1)}{r^2}\varphi_l = 0, \tag{11}$$

where we have separated out the angular part of $\phi(x)$ in Eq. (7). There is no exact solution to this equation. We therefore adopt a perturbative approach assuming that the quantity,

$$\epsilon \equiv GMH, \tag{12}$$

is a small parameter. But the condition (5) implies that

$$\epsilon > 3.96 \times 10^{-2}\left(\frac{H}{M_{\rm Pl}}\right)^{\frac{2}{3}}. \tag{13}$$

However, this shows that the smallness of ϵ can be easily satisfied for any reasonable inflation scale. Let us rewrite Eq. (11) as

$$\begin{aligned}\partial_\tau^2\varphi_l - \frac{2}{\tau}\partial_\tau\varphi_l - \partial_r^2\varphi_l - \frac{2}{r}\partial_r\varphi_l + \frac{l(l+1)}{r^2}\varphi_l &= \left(1-\frac{h}{f}\right)\partial_\tau^2\varphi_l - \frac{2}{\tau}\left(1-\frac{h}{f}\right)\partial_\tau\varphi_l \\ &-\frac{1}{\tau^2\sqrt{fh}}\partial_\tau\left(\tau^2\sqrt{\frac{h^3}{f}}\right)\partial_\tau\varphi_l + \frac{1}{\sqrt{fh}}\partial_r\left(\sqrt{fh}\right)\partial_r\varphi_l\end{aligned} \tag{14}$$

We then expand the functions f and h in powers of ϵ as defined in Eq. (12) and write

$$\varphi_l = \varphi_l^{(0)} + \varphi_l^{(1)} + \varphi_l^{(2)} + \cdots \tag{15}$$

as an expansion in orders of ϵ. It is straightforward to show that the right-hand side of Eq. (14) becomes

$$8\left(\frac{\epsilon\tau}{2r}\right)\left(\partial_\tau^2\varphi_l - \frac{1}{\tau}\partial_\tau\varphi_l\right) - 30\left(\frac{\epsilon\tau}{2r}\right)^2\left(\partial_\tau^2\varphi_l - \frac{1}{15\tau}\partial_\tau\varphi_l - \frac{1}{15r}\partial_r\varphi_l\right) + \cdots \tag{16}$$

1. Zeroth order

From Eqs. (14), (15), and (16), the zeroth order corresponds to the de Sitter case with

$$\partial_\tau^2\varphi_l^{(0)} - \frac{2}{\tau}\partial_\tau\varphi_l^{(0)} - \partial_r^2\varphi_l^{(0)} - \frac{2}{r}\partial_r\varphi_l^{(0)} + \frac{l(l+1)}{r^2}\varphi_l^{(0)} = 0. \tag{17}$$

To solve this equation, we take the Bessel transform,

$$\varphi_l^{(0)}(r,\tau) = \int_0^\infty dk k^2 j_l(kr)\varphi_{kl}^{(0)}(\tau) \ \ . \tag{18}$$

Then we have

$$\partial_\tau^2\varphi_{kl}^{(0)}(\tau) - \frac{2}{\tau}\partial_\tau\varphi_{kl}^{(0)}(\tau) + k^2\varphi_{kl}^{(0)}(\tau) = 0 \ \ , \tag{19}$$

and the solution is

$$\varphi_{kl}^{(0)}(\tau) = C_1(-k\tau)^{\frac{3}{2}}H_{\frac{3}{2}}^{(1)}(-k\tau) + C_2(-k\tau)^{\frac{3}{2}}H_{\frac{3}{2}}^{(2)}(-k\tau) \ . \tag{20}$$

If we take the boundary conditions:

$$C_1 = -\frac{H}{k^2\sqrt{2k}}, \quad \text{and} \quad C_2 = 0, \tag{21}$$

then we will have

$$\varphi_{kl}^{(0)}(\tau) = -\frac{H\tau}{k\sqrt{\pi k}}\left(1 - \frac{i}{k\tau}\right)e^{-ik\tau}, \tag{22}$$

and

$$P_{kl}^{(0)}(\tau) = \frac{k^5}{4\pi}\left|\varphi_{kl}^{(0)}(\tau)\right|^2 = \frac{H^2}{4\pi^2}(1 + k^2\tau^2). \tag{23}$$

As $\tau \to 0$, $P_{kl}^{(0)}(\tau) \to H^2/(4\pi^2)$. This result matches the well-known scale-invariant power spectrum of de Sitter quantum fluctuations [10].

2. First order

With the perturbative expansion for φ_l in Eq. (15), the Klein-Gordon equation in Eq. (14) can be solved perturbatively. The first order $\varphi_l^{(1)}(r,\tau)$ then satisfies

$$\partial_\tau^2\varphi_l^{(1)} - \frac{2}{\tau}\partial_\tau\varphi_l^{(1)} - \partial_r^2\varphi_l^{(1)} - \frac{2}{r}\partial_r\varphi_l^{(1)} + \frac{l(l+1)}{r^2}\varphi_l^{(1)} = J_1 \ , \tag{24}$$

where the source term is given by

$$\begin{aligned} J_1(r,\tau) &= \frac{4\epsilon\tau}{r}\left(\partial_\tau^2\varphi_l^{(0)} - \frac{1}{\tau}\partial_\tau\varphi_l^{(0)}\right) \\ &= \frac{4\epsilon H\tau^2}{\sqrt{\pi}r}\int_0^\infty dk k^2 j_l(kr)(k^{\frac{1}{2}}e^{-ik\tau}). \end{aligned} \tag{25}$$

To solve this inhomogeneous equation, we use the Green's function $G(r,\tau;r',\tau')$ which satisfies the equation,

$$\partial_\tau^2 G - \frac{2}{\tau}\partial_\tau G - \partial_r^2 G - \frac{2}{r}\partial_r G + \frac{l(l+1)}{r^2}G = \frac{\delta(r-r')\delta(\tau-\tau')}{r^2} \ . \tag{26}$$

Using the completeness property of the spherical Bessel functions,

$$\int_0^\infty dk k^2 \left[\sqrt{\frac{2}{\pi}} r j_l(kr)\right]\left[\sqrt{\frac{2}{\pi}} r' j_l(kr')\right] = \delta(r-r'), \tag{27}$$

and taking

$$G_l(r,\tau;r',\tau') = \int_0^\infty dk k^2 g_k(\tau,\tau') j_l(kr) j_l(kr') \ , \tag{28}$$

Eq. (26) becomes

$$\partial_\tau^2 g_k - \frac{2}{\tau}\partial_\tau g_k + k^2 g_k = \frac{2}{\pi}\delta(\tau-\tau') \ . \tag{29}$$

For the retarded Green's function, $g_k = 0$ for $\tau' > \tau > \tau_i$, where τ_i denotes an initial time when the source begins to operate. For $0 > \tau > \tau'$,

$$\begin{aligned} g_k(\tau,\tau') \ &= \ \frac{i}{2\tau'^2 k^3}\Big[(-k\tau)^{\frac{3}{2}} H^{(1)}_{\frac{3}{2}}(-k\tau)(-k\tau')^{\frac{3}{2}} H^{(2)}_{\frac{3}{2}}(-k\tau') \\ &\qquad - (-k\tau')^{\frac{3}{2}} H^{(1)}_{\frac{3}{2}}(-k\tau')(-k\tau)^{\frac{3}{2}} H^{(2)}_{\frac{3}{2}}(-k\tau)\Big] \ . \end{aligned} \tag{30}$$

With this retarded Green's function, the first order $\varphi_l^{(1)}(r,\tau)$ can be expressed as

$$\varphi_l^{(1)}(r,\tau) = \int_0^\infty dk k^2 j_l(kr)\varphi_{kl}^{(1)}(\tau) = \int_0^\infty dr' r'^2 \int_{\tau_i}^0 d\tau' G(r,\tau;r',\tau') J_1(r',\tau'). \tag{31}$$

Hence, we find that

$$\begin{aligned} \varphi_{kl}^{(1)}(\tau) \ &= \ \frac{2i\epsilon H}{\sqrt{\pi}k^3}\int_0^\infty dk' k'^2 (k'^{\frac{1}{2}}) \int_0^\infty dr' r' j_l(kr') j_l(k'r') \int_{\tau_i}^{\tau} d\tau' e^{-ik'\tau'} \times \\ &\Big[(-k\tau)^{\frac{3}{2}} H^{(1)}_{\frac{3}{2}}(-k\tau)(-k\tau')^{\frac{3}{2}} H^{(2)}_{\frac{3}{2}}(-k\tau') - (-k\tau')^{\frac{3}{2}} H^{(1)}_{\frac{3}{2}}(-k\tau')(-k\tau)^{\frac{3}{2}} H^{(2)}_{\frac{3}{2}}(-k\tau)\Big] \ . \end{aligned} \tag{32}$$

It is useful to rewrite $\varphi_{kl}^{(1)}(\tau)$ as

$$\varphi_{kl}^{(1)}(\tau) = \epsilon\left[\alpha_{kl}(\tau)\varphi_{kl}^{(0)}(\tau) + \beta_{kl}(\tau)\varphi_{kl}^{(0)*}(\tau)\right]. \tag{33}$$

The integral over r' can be performed,

$$\int_0^\infty dr' r' j_l(kr') j_l(k'r') = \frac{\pi}{2\sqrt{k_< k_>}}\left(\frac{1}{k_>}\right)\left(\frac{k_<}{k_>}\right)^{l+\frac{1}{2}} \frac{\Gamma(l+1)}{\Gamma(l+\frac{3}{2})\Gamma(\frac{1}{2})} F\left(l+1,\frac{1}{2};l+\frac{3}{2},\frac{k_<^2}{k_>^2}\right) \ , \tag{34}$$

and after some rescalings the coefficients α_{kl} and β_{kl} can be simplified to

$$\begin{aligned}\alpha_{kl}(\tau) &= \frac{-2i\Gamma(l+1)}{\Gamma(l+\frac{3}{2})\Gamma(\frac{1}{2})}\left[\int_0^1 dk' k'^{l+\frac{5}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};k'^2\right)\int_{k\tau_i}^{k\tau} d\tau' e^{-ik'\tau'} e^{i\tau'}\left(\tau'+i\right)\right.\\ &\quad \left. + \int_1^\infty dk' k'^{-l+\frac{1}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};\frac{1}{k'^2}\right)\int_{k\tau_i}^{k\tau} d\tau' e^{-ik'\tau'} e^{i\tau'}\left(\tau'+i\right)\right] \qquad (35)\end{aligned}$$

and

$$\begin{aligned}\beta_{kl}(\tau) &= \frac{2i\Gamma(l+1)}{\Gamma(l+\frac{3}{2})\Gamma(\frac{1}{2})}\left[\int_0^1 dk' k'^{l+\frac{5}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};k'^2\right)\int_{k\tau_i}^{k\tau} d\tau' e^{-ik'\tau'} e^{-i\tau'}\left(\tau'-i\right)\right.\\ &\quad \left. + \int_1^\infty dk' k'^{-l+\frac{1}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};\frac{1}{k'^2}\right)\int_{k\tau_i}^{k\tau} d\tau' e^{-ik'\tau'} e^{-i\tau'}\left(\tau'-i\right)\right]. \qquad (36)\end{aligned}$$

The power spectrum in this order is then given by

$$P_{kl}(\tau) = P_{kl}^{(0)}(\tau)\left[1+\epsilon\Delta_{kl}^{(1)}(\tau)\right], \qquad (37)$$

where we have introduced the dimensionless quantity,

$$\begin{aligned}\Delta_{kl}^{(1)}(\tau) &= 2\,\epsilon^{-1}\left|\varphi_{kl}^{(0)}(\tau)\right|^{-2}\mathrm{Re}\left[\varphi_{kl}^{(0)}(\tau)\varphi_{kl}^{(1)*}(\tau)\right]\\ &= 2\,\mathrm{Re}\left[\alpha_{kl}(\tau)\right] + 2\left|\varphi_{kl}^{(0)}(\tau)\right|^{-2}\mathrm{Re}\left[\beta_{kl}^{*}(\tau)\varphi_{kl}^{(0)2}(\tau)\right]. \qquad (38)\end{aligned}$$

As $\tau \to 0$, from Eq. (22),

$$\Delta_{kl}^{(1)}(0) = 2\,\mathrm{Re}\left[\alpha_{kl}(0)-\beta_{kl}(0)\right]. \qquad (39)$$

From Eqs. (35) and (36), we find that

$$\begin{aligned}\Delta_{kl}^{(1)}(0) &= \frac{8\Gamma(l+1)}{\Gamma(l+\frac{3}{2})\Gamma(\frac{1}{2})}\left[\int_0^1 dk' k'^{l+\frac{5}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};k'^2\right)+\right.\\ &\quad \left.\int_1^\infty dk' k'^{-l+\frac{1}{2}} F\left(l+1,\frac{1}{2};l+\frac{3}{2};\frac{1}{k'^2}\right)\right]\times\\ &\quad \frac{1}{(k'^2-1)^2}\left\{\cos(k\tau_i)\left[2\sin(k'k\tau_i)-k'(k'^2-1)k\tau_i\cos(k'k\tau_i)\right]\right.\\ &\quad \left.+\sin(k\tau_i)\left[k'(k'^2-3)\cos(k'k\tau_i)-(k'^2-1)k\tau_i\sin(k'k\tau_i)\right]\right\}. \qquad (40)\end{aligned}$$

B. Quantization

A unique mode function $\varphi_{klm}(x)$ can be obtained once an appropriate vacuum is chosen. By using this mode function the scalar field can be quantized in the standard manner,

$$\hat{\phi}(x) = \int_0^\infty dk \sum_{lm}\left[\hat{a}_{klm}\varphi_{klm}(x)+\hat{a}_{klm}^{\dagger}\varphi_{klm}^{*}(x)\right], \qquad (41)$$

with the commutation relations:

$$\begin{aligned}
&[\hat{a}_{klm}, \hat{a}_{k'l'm'}] = [\hat{a}^\dagger_{klm}, \hat{a}^\dagger_{k'l'm'}] = 0, \\
&[\hat{a}_{klm}, \hat{a}^\dagger_{k'l'm'}] = \delta(k-k')\delta_{ll'}\delta_{mm'}.
\end{aligned} \tag{42}$$

We have the delta function in k because there is no coupling between different k-modes. We will show this explicitly later by a perturbation approach. The delta functions in l and m stem from the rotational invariance about the central black hole. The vacuum state is defined as

$$\hat{a}_{klm}|0\rangle = 0. \tag{43}$$

Now $\varphi_{klm}(x)$ is the mode function, so it should also satisfy the Klein-Gordon equation. Since the space is spherically symmetric, one can write

$$\varphi_{klm}(x) = \varphi_{kl}(r,\tau) Y_{lm}(\theta,\phi). \tag{44}$$

But $\varphi_{kl}(r,\tau)$ cannot in general be separated as a product of functions with only one variable like $k^2 j_l(kr)\varphi_{kl}(\tau)$ as we have done in the classical solution. However, $\varphi_{kl}(r,\tau)$ can still be obtained perturbatively. The Klein-Gordon equation for the mode function is the same as Eq. (11), given by

$$-\frac{1}{\sqrt{fh}}\partial_\tau\left(\sqrt{\frac{h^3}{f}}\partial_\tau\varphi_{kl}(r,\tau)\right) + \frac{1}{r^2\sqrt{fh}}\partial_r\left(r^2\sqrt{fh}\partial_r\varphi_{kl}(r,\tau)\right) - \frac{l(l+1)}{r^2}\varphi_{kl}(r,\tau) = 0. \tag{45}$$

The two-point correlation function is then given by

$$\begin{aligned}
\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle &= \int_0^\infty dk \sum_{lm} \varphi_{klm}(x)\varphi^*_{klm}(x') \\
&= \int_0^\infty dk \sum_l \frac{2l+1}{4\pi}\varphi_{kl}(r,\tau)\varphi^*_{kl}(r',\tau)P_l(\cos\gamma),
\end{aligned} \tag{46}$$

where γ is the separation angle between the two points. As $x' \to x$, we have

$$\langle 0|\hat{\phi}^2(x)|0\rangle = \int_0^\infty \frac{dk}{k}\sum_l \frac{2l+1}{4\pi} k\left|\varphi_{kl}(r,\tau)\right|^2. \tag{47}$$

In terms of $\varphi_{kl}(r,\tau)$, the spectral function of the fluctuations of the quantum field $\phi(x)$ can be defined as

$$S_{kl}(r,\tau) \equiv (2l+1)j_l^2(kr)P_{kl}(r,\tau), \quad P_{kl}(r,\tau) = \frac{k}{4\pi j_l^2(kr)}\left|\varphi_{kl}(r,\tau)\right|^2. \tag{48}$$

Perturbatively,

$$\begin{aligned}
P_{kl}(r,\tau) &= \frac{k}{4\pi j_l^2(kr)}\left|\varphi^{(0)}_{kl}(r,\tau) + \varphi^{(1)}_{kl}(r,\tau) + \cdots\right|^2 \\
&= P^{(0)}_{kl}(r,\tau)\left[1 + \epsilon\Delta^{(1)}_{kl}(r,\tau) + \cdots\right],
\end{aligned} \tag{49}$$

where we have defined

$$P^{(0)}_{kl}(r,\tau) = \frac{k}{4\pi j_l^2(kr)}\left|\varphi^{(0)}_{kl}(r,\tau)\right|^2, \tag{50}$$

$$\Delta^{(1)}_{kl}(r,\tau) = 2\,\epsilon^{-1}\left|\varphi^{(0)}_{kl}(r,\tau)\right|^{-2}\mathrm{Re}\left[\varphi^{(0)}_{kl}(r,\tau)\varphi^{(1)*}_{kl}(r,\tau)\right]. \tag{51}$$

1. *Zeroth order*

Following the same steps in Eqs. (14), (15), and (16), to the zeroth order we have as before

$$\partial_\tau^2\varphi_{kl}^{(0)}(r,\tau)-\frac{2}{\tau}\partial_\tau\varphi_{kl}^{(0)}(r,\tau)-\partial_r^2\varphi_{kl}^{(0)}(r,\tau)-\frac{2}{r}\partial_r\varphi_{kl}^{(0)}(r,\tau)+\frac{l(l+1)}{r^2}\varphi_{kl}^{(0)}(r,\tau)=0, \tag{52}$$

and the general solution is found to be

$$\varphi_{kl}^{(0)}(r,\tau)=k^2 j_l(kr)\varphi_{kl}^{(0)}(\tau), \tag{53}$$

where $\varphi_{kl}^{(0)}(\tau)$ is given by Eq. (20). The boundary conditions (21) indeed correspond to the choice of the Bunch-Davies vacuum that selects the mode function:

$$\varphi_{kl}^{(0)}(\tau)=-\frac{H\tau}{k\sqrt{\pi k}}\left(1-\frac{i}{k\tau}\right)e^{-ik\tau}, \tag{54}$$

and hence the zeroth order power spectrum in Eq. (50) is

$$P_{kl}^{(0)}(r,\tau)=\frac{k}{4\pi j_l^2(kr)}\left|\varphi_{kl}^{(0)}(r,\tau)\right|^2=\frac{H^2}{4\pi^2}(1+k^2\tau^2). \tag{55}$$

2. *First order*

To the next order, we have

$$\partial_\tau^2\varphi_{kl}^{(1)}(r,\tau)-\frac{2}{\tau}\partial_\tau\varphi_{kl}^{(1)}(r,\tau)-\partial_r^2\varphi_{kl}^{(1)}(r,\tau)-\frac{2}{r}\partial_r\varphi_{kl}^{(1)}(r,\tau)+\frac{l(l+1)}{r^2}\varphi_{kl}^{(1)}(r,\tau)=J_1\ , \tag{56}$$

where

$$\begin{aligned}J_1(r,\tau)\ &=\ \frac{4\epsilon\tau}{r}\left[\partial_\tau^2\varphi_{kl}^{(0)}(r,\tau)-\frac{1}{\tau}\partial_\tau\varphi_{kl}^{(0)}(r,\tau)\right]\\ &=\ \frac{4\epsilon H\tau^2}{\sqrt{\pi}r}\left[k^2 j_l(kr)(k^{\frac{1}{2}}e^{-ik\tau})\right].\end{aligned} \tag{57}$$

The retarded Green's function necessary to solve this equation is the same as before. Then,

$$\begin{aligned}&\varphi_{kl}^{(1)}(r,\tau)\\ &=\int_0^\infty dr'r'^2\int_{\tau_i}^0 d\tau' G(r,\tau;r',\tau')J_1(r',\tau')\\ &=\frac{2i\epsilon H\ k^{\frac{5}{2}}}{\sqrt{\pi}}\int_0^\infty\frac{dk'}{k'}j_l(k'r)\int_0^\infty dr'r'j_l(kr')j_l(k'r')\int_{\tau_i}^\tau d\tau' e^{-ik\tau'}\times\\ &\left[(-k'\tau)^{\frac{3}{2}}H_{\frac{3}{2}}^{(1)}(-k'\tau)(-k'\tau')^{\frac{3}{2}}H_{\frac{3}{2}}^{(2)}(-k'\tau')-(-k'\tau')^{\frac{3}{2}}H_{\frac{3}{2}}^{(1)}(-k'\tau')(-k'\tau)^{\frac{3}{2}}H_{\frac{3}{2}}^{(2)}(-k'\tau)\right].\end{aligned} \tag{58}$$

This is in the form of

$$\varphi^{(1)}_{kl}(r,\tau) = \epsilon \int_0^\infty \frac{dk'}{k'} \left[\alpha_{kk'l}(\tau)\varphi^{(0)}_{k'l}(r,\tau) + \beta_{kk'l}(\tau){\varphi^{(0)}_{k'l}}^*(r,\tau)\right], \tag{59}$$

where

$$\begin{aligned} \alpha_{kk'l}(\tau) &= -\left(\frac{4i}{\pi}\right) k^{\frac{5}{2}} k'^{\frac{1}{2}} \int_0^\infty dr'\, r' j_l(kr') j_l(k'r') \int_{\tau_i}^{\tau} d\tau' (k'\tau' + i) e^{-i(k-k')\tau'}, &(60)\\ \beta_{kk'l}(\tau) &= \left(\frac{4i}{\pi}\right) k^{\frac{5}{2}} k'^{\frac{1}{2}} \int_0^\infty dr'\, r' j_l(kr') j_l(k'r') \int_{\tau_i}^{\tau} d\tau' (k'\tau' - i) e^{-i(k+k')\tau'}. &(61)\end{aligned}$$

As the perturbative formalism in quantum mechanics, the eigenfunctions in the first order consist of zeroth order eigenfunctions of different energies.

Now we calculate the leading correction (51) to the power spectrum. Substituting Eq. (53) in Eq. (59), we have

$$\Delta^{(1)}_{kl}(r,\tau) = \int_0^\infty \frac{dk'}{k'}\frac{k'^2}{k^2}\frac{j_l(k'r)}{j_l(kr)} \left|\varphi^{(0)}_{kl}(\tau)\right|^{-2} 2\,\mathrm{Re}\left[\alpha^*_{kk'l}(\tau)\varphi^{(0)}_{kl}(\tau)\varphi^{(0)*}_{k'l}(\tau) + \beta^*_{kk'l}(\tau)\varphi^{(0)}_{kl}(\tau)\varphi^{(0)}_{k'l}(\tau)\right]. \tag{62}$$

As $\tau \to 0$, this becomes

$$\Delta^{(1)}_{kl}(r,0) = \int_0^\infty \frac{dk'}{k'}\left(\frac{k}{k'}\right)^{\frac{1}{2}} \frac{j_l(k'r)}{j_l(kr)}\, 2\,\mathrm{Re}\left[\alpha_{kk'l}(0) - \beta_{kk'l}(0)\right]. \tag{63}$$

Here we obtain from Eqs. (60) and (61) that

$$\begin{aligned} & 2\,\mathrm{Re}\left[\alpha_{kk'l}(0) - \beta_{kk'l}(0)\right] \\ = \; & \frac{16}{\pi}\left(\frac{k}{k'}\right)^{\frac{1}{2}} k^2 \int_0^\infty dr'\, r' j_l(kr') j_l(k'r') \times \\ & \left(\frac{k^2}{k'^2} - 1\right)^{-2} \left\{ \cos(k'\tau_i)\left[2\sin(k\tau_i) - \left(\frac{k^2}{k'^2} - 1\right) k\tau_i \cos(k\tau_i)\right] \right. \\ & \left. + \sin(k'\tau_i)\left[\frac{k}{k'}\left(\frac{k^2}{k'^2} - 3\right)\cos(k\tau_i) - \left(\frac{k^2}{k'^2} - 1\right) k'\tau_i \sin(k\tau_i)\right]\right\}, \end{aligned} \tag{64}$$

where the remaining integral over r' can be evaluated using the formula (34). This quantity diverges as $k \to k'$. However, we find that the integral in Eq. (63) is finite and weakly dependent on r. In the limit of $r \to 0$, it can be approximated by

$$\Delta^{(1)}_{kl}(r,0) \simeq \int_0^\infty \frac{dk'}{k'}\left(\frac{k'}{k}\right)^{l-\frac{1}{2}} 2\,\mathrm{Re}\left[\alpha_{kk'l}(0) - \beta_{kk'l}(0)\right]. \tag{65}$$

III. NUMERICAL RESULTS

Eqs. (38) and (62) are the main results of the present paper. However, they are complicated integrals and are not illuminating. Therefore, we perform numerical calculations of the black-hole corrections to the power spectrum. Assume that inflation begins at the initial time $t = 0$ and $a(t = 0) = 1$. Then, the initial conformal time is $\tau_i = -1/H$ and the final conformal time is $\tau = -e^{-Ht}/H$, where Ht is the e-folding number of inflation. It is useful to note that the Fourier mode with momentum k crosses out the cosmological horizon at time $\tau = -1/k$, or equivalently, when the e-folding number is $Ht = \ln(k/H)$.

In Fig. 1, we show the time evolution of the first-order contribution to the quantum fluctuations from Eq. (37). Actually, it is more convenient to plot the normalized power spectrum $\Delta^{(1)}_{kl}(\tau)$ in Eq. (38) against the e-folding number Ht. We have chosen the angular momentum $l = 2, 22, 40$ as examples. For each l, different k modes are shown. From all the plots, we can see that for a k-mode $\Delta^{(1)}_{kl}(\tau)$ oscillates when the mode is still sub-horizon. Once the mode crosses out the horizon, $\Delta^{(1)}_{kl}(\tau)$ stop oscillating and gradually approaches a constant value. This behavior can be easily explained by Eq. (25), where the source term dies off as $\varphi^{(0)}_{kl}$ goes super-horizon and then gets frozen. In Fig. 2, we plot the asymptotic values $\Delta^{(1)}_{kl}(0)$ in Eq. (39) for $1 < l < 50$ and $1 < k/H < 50$. The figure shows that $\Delta^{(1)}_{kl}(0)$ is suppressed in low-l and low-k regions and the magnitude of $|\Delta^{(1)}_{kl}(0)|$ is of order one. Thus, the first-order contribution, compared to the zero-order de Sitter power spectrum, is roughly downsized by the expansion parameter ϵ.

For the quantum case, we plot the asymptotic values $\Delta^{(1)}_{kl}(r, 0)$ in Eq. (63) for $Hr = 0.01, 0.1$ against l and k/H for $1 < l < 50$ and $1 < k/H < 50$. As shown in Fig. 3, the general trend is similar to the classical case in Fig. 2. This is indeed an explicit example that quantum fluctuations generated during inflation behave like classical waves. We have also calculated $\Delta^{(1)}_{kl}(r, 0)$ for larger values of Hr, which becomes fluctuating but in general the value of the amplitude is getting smaller. It is expected because the farther the black hole is, the lesser is its effect to the perturbation.

IV. CONCLUSIONS

We have presented a perturbation method to compute the effect of the presence of a black hole in the de Sitter space to the quantum fluctuations of a free massless scalar field. The method is valid as long as the expansion parameter $\epsilon \equiv GMH < 1$, i.e., the product of the mass of the black hole and the de Sitter Hubble parameter has to be smaller than the Planck mass squared. The calculation can be easily applied to a vector field or a gravitational wave. Here the first-order contribution is computed and the results are given in the assumption that the black hole is located at the origin of the coordinates. Higher-order corrections can be worked out perturbatively though complicated. It would be interesting to consider the effect due to a distribution of black holes in the de Sitter space. In fact, the perturbation can be in a form of cosmological defects such as monopoles, cosmic strings, or domain walls.

Let us briefly discuss some cosmological implications of the results that we have obtained in this work. Assume that inflation lasts for about 60 e-folds. Then, the wavelength of the Fourier mode with $k/H = 1$ is about the size of the present Universe. If the location of the black hole that exists during inflation is near the Earth, the suppressed power of the first-order correction to the de Sitter inflaton fluctuations in low l and low k regions may result in a blue-tilted density power spectrum on large angular scales. This in turn gives rise to a suppression of the large-scale cosmic microwave background anisotropy that may be relevant to the observed low quadrupole in the WMAP cosmic microwave background anisotropy data [11]. A detailed calculation of the effect to the cosmic microwave background anisotropy is underway, including the case that the black hole locates somewhere else in the

Universe.

If inflation lasts for a longer time, then the wavelength of the Fourier mode that corresponds to the size of the present Universe will be given by a larger value of k/H. In Fig. 4, we plot the asymptotic values $\Delta_{kl}^{(1)}(0)$ against l for $k/H = 1$, 10, 100, and 250, which correspond to the inflation with 60, 62.3, 64.6, and 65.5 e-folds, respectively. As expected, the longer is the inflation duration the lesser pronounced are the effects of the black hole to large-scale or low-l observations. This can also be seen in the quantum case with $\Delta_{kl}^{(1)}(r, 0)$ in Fig. 3.

It is worth noting that the present work gives a realization of the general discussions in Ref. [12] about the potentially observable effects of a small violation of translational invariance during inflation, as characterized by the presence of a preferred point, line, or plane. This violation may induce derivations from pure statistical isotropy of cosmological perturbations, thus leaving anomalous imprints on the cosmic microwave background anisotropy [12].

Acknowledgments

We thank Lau-Loi So for his contributions at the initial stage of the paper and the anonymous referee for his/her comments. This work was supported in part by the National Science Council, Taiwan, ROC under the Grants NSC 96-2112-M-032-006-MY3 (HTC), NSC 95-2112-M-001-052-MY3 (KWN), NSC 97-2112-M-003-004-MY3 (ICW), and the National Center for Theoretical Sciences, Taiwan, ROC.

---

[1] E. Komatsu *et al.*, Astrophys. J. Suppl. **180**, 330 (2009).
[2] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2738 (1977).
[3] H. Kurki-Suonio, J. Centrella, R. A. Matzner, and J. R. Wilson, Phys. Rev. D **35**, 435 (1987); D. S. Goldwirth and T. Piran, Phys. Rev. D **40**, 3263 (1989); P. Laguna, H. Kurki-Suonio, and R. A. Matzner, Phys. Rev. D **44**, 3077 (1991).
[4] D. S. Goldwirth and T. Piran, Phys. Rev. Lett. **64**, 2852 (1990); E. Calzetta and M. Sakellariadou, Phys. Rev. D **45**, 2802 (1992).
[5] K. A. Holcomb, S. J. Park, and E. T. Vishniac, Phys. Rev. D **39**, 1058 (1989).
[6] D. Garfinkle and C. Vuille, Gen. Relativ. Gravit. **23**, 471 (1991); K. Nakao, Gen. Relativ. Gravit. **24**, 1069 (1992).
[7] K. Nakao, K. Maeda, T. Nakamura, and K. Oohara, Phys. Rev. D **47**, 3194 (1993).
[8] G. C. MacVittie, Mon. Not. Roy. Astron. Soc. **93**, 325 (1933); M. Kihara and H. Nariai, Prog. Theor. Phys. **65**, 1613 (1981); T. Shiromizu, D. Ida, and T. Torii, J. High Energy Phys. **11**, 010 (2001).
[9] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975); Erratum-ibid. **46**, 206 (1976).
[10] S. W. Hawking, Phys. Lett. B **115**, 295 (1982); A. A. Starobinsky, Phys. Lett. B **117**, 175 (1982); A. H. Guth and S. Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
[11] M. R. Nolta *et al.*, Astrophys. J. Suppl. **180**, 296 (2009).
[12] S. M. Carroll, C.-Y. Tseng, and M. B. Wise, arXiv:0811.1086.



FIG. 1: Time evolution of the first-order normalized power spectra $\Delta_{kl}^{(1)}(\tau)$ in Eq. (38) for $l = 2, 22, 40$. For each case, we have selected some k-modes. The time is expressed in terms of the e-folding number Ht.



FIG. 2: Two views of the three-dimensional plot of the asymptotic values $\Delta_{kl}^{(1)}(0)$ in Eq. (39) against l and k/H for $1 < l < 50$ and $1 < k/H < 50$.





FIG. 3: Three-dimensional plots of the asymptotic values $\Delta_{kl}^{(1)}(r,0)$ in Eq. (63) for $Hr = 0.01$, 0.1 against l and k/H for $1 < l < 50$ and $1 < k/H < 50$.



FIG. 4: Asymptotic values $\Delta_{kl}^{(1)}(0)$ against l for $k/H = 1$, 10, 100, and 250.